

S.E. BTRX/EXTC - Applied Mathematics - III

CBGS

Q.P. Code : 13607

16/05/2018

(3 Hours)

[Total marks : 80]

- Note :-
- 1) Question number 1 is compulsory.
 - 2) Attempt any three questions from the remaining five questions.
 - 3) Figures to the right indicate full marks.

- Q.1
- Find the Laplace transform of $\cos t \cos 2t \cos 3t$. 05
 - Construct an analytic function whose real part is $e^x \cos y$. 05
 - Find the directional derivative of $\phi = x^2 + y^2 + z^2$ at point $A(1, -2, 1)$ in the direction of AB where B is $(2, 6, -1)$. 05
 - Expand $f(x) = lx - x^2$, $0 < x < l$ in a half-range sine-series. 05
- Q.2
- Find the angle between the normals to the surface $xy = z^2$ at the points $(1, 4, 2)$, $(-3, -3, 3)$. 06
 - Find the Fourier series for $f(x) = \begin{cases} -c & -a < x < 0 \\ c & 0 < x < a \end{cases}$ 06
 - Find the inverse Laplace transform of $\frac{4s + 12}{s^2 + 8s + 12}$ 08
- (i) $\frac{4s + 12}{s^2 + 8s + 12}$
- (ii) $\log\left(\frac{s^2 + a^2}{\sqrt{s + b}}\right)$
- Q.3
- State true or false with proper justification "There does not exist an analytic function whose real part is $x^3 - 3x^2y - y^3$ ". 06
 - Prove that $\int_{5/2}^x \sqrt{2} \left(\frac{3-x^2}{x^2} \sin x - \frac{3}{x} \cos x \right) dx = \frac{2}{\pi x} \left(\frac{3-x^2}{x^2} \sin x - \frac{3}{x} \cos x \right)$. 06
 - Expand $f(x) = 4 - x^2$ in the interval $(0, 2)$. 08
- Q.4
- Use Gauss's Divergence theorem to evaluate $\iint_S \vec{N} \cdot \vec{F} dS$ where $\vec{F} = 4x\vec{i} + 3y\vec{j} - 2z\vec{k}$ and S is the surface bounded by $x = 0, y = 0, z = 0$ and $2x + 2y + z = 4$. 06

TURN OVER

- b) Prove that $\int x^3 \cdot J_0(x) dx = x^3 \cdot J_1(x) - 2x^2 \cdot J_2(x)$. 06
- c) Solve using Laplace transform $\frac{dy}{dt} + 3y = 2 + e^{-t}$ with $y(0) = 1$. 08

Q. 5 a) Find Laplace transform of $(1 + 2t - 3t^2 + 4t^3)H(t - 2)$ where $H(t - 2) = \begin{cases} 0, & t < 2 \\ 1, & t \geq 2 \end{cases}$ 06

- b) Prove that $2J_0''(x) = J_2(x) - J_0(x)$. 06
- c) Obtain complex form of Fourier Series for $f(x) = e^{ax}$ in $(-\pi, \pi)$ where a is not an integer. Hence deduce that when a is a constant other than an integer 08

$$\sin ax = \frac{\sin \pi a}{i\pi} \sum \frac{(-1)^n n}{(a^2 - n^2)} e^{inx}$$

Q. 6 a) Using Green's theorem evaluate 06

$$\oint_C (e^{x^2} - xy) dx - (y^2 - ax) dy$$

where C is the circle $x^2 + y^2 = a^2$.

- b) Express the function $f(x) = \begin{cases} 1 & \text{for } |x| < 1 \\ 0 & \text{for } |x| > 1 \end{cases}$ as a Fourier Integral. 06
- c) Under the transformation $w = (1 + i)z + (2 - i)$, find the region in the w -plane into which the rectangular region bounded by $x = 0, y = 0, x = 1, y = 2$ in the z -plane is mapped. 08

xxx xxx